

## **HEALTHY ALASKANS 2020**

A JOINT PROJECT OF THE STATE OF ALASKA DEPARTMENT OF HEALTH AND SOCIAL SERVICES & THE ALASKA NATIVE TRIBAL HEALTH CONSORTIUM

## Healthy Alaskans 2020 Scorecard **Regional Edition - Mat-Su**

	HA2020 Leading Health Indicator	Mat-Su Baseline*	HA2020 Target	Mat-Su Current*	Target Met?	
1	Reduce the cancer mortality rate per 100,000 population	201.4	162.0	144.7	$\checkmark$	
2	Increase the percentage of adolescents (high school students in grades 9-12) who have not smoked cigarettes or cigars or used chew-ing tobacco, snuff or dip on one or more of the past 30 days	74.0%ª	80%	78.5%	Х	
3	Increase the percentage of adults (age 18 and older) who do not currently smoke cigarettes	71.0%	83%	83.3%	$\checkmark$	
4a	Reduce the percentage of adults (age 18 years and older) who meet criteria for overweight (body mass index of ≥ 25 and < 30 kg/m <sup>2</sup> )	38.6%	36%	35.6%	$\checkmark$	
4b	Reduce the percentage of adults (age 18 years and older) who meet criteria for obesity (body mass index of ≥ 30 kg/m <sup>2</sup> )	37.5%	27%	32.8%	Х	
5ai	Reduce the percentage of adolescents (high school students in grades 9-12) who meet criteria for overweight (age- and sex-specific body mass index of ≥ 85th and < 95th percentile)	11.2%ª	12%	16.5%	Х	
5aii	Reduce the percentage of adolescents (high school students in grades 9-12) who meet criteria for obesity (age- and sex-specific body mass index of $\geq$ 95th percentile)	12.0%ª	10%	14.3%	Х	
5bi	Reduce the percentage of children (students in grades K-8) who meet criteria for overweight (age- and sex-specific body mass index of >=85th and <95th percentile)	Statewide Only	15%	Statewide Only		
5bii	Reduce the percentage of children (students in grades K-8) who meet criteria for obesity (age- and sex-specific body mass index of >=95th percentile)	Statewide Only	15%	Statewide Only		
6a	Increase the percentage of adults (age 18 years and older) who report 150 or more minutes per week of moderate or vigorous exer- cise, where each minute of vigorous exercise contributes 2 minutes to the total	55.9% <sup>b</sup>	61%	58.9%	Х	
6b	Increase the percentage of adolescents (high school students in grades 9-12) who do at least 60 minutes of physical activity a day, every day of the week	19.3%ª	23%	19.4%	Х	
7a	Reduce the suicide mortality rate per 100,000 population, among the population age 15-24 years	DSU	43.2	DSU		
7b	Reduce the suicide mortality rate per 100,000 population, among the population age 25 years and older	26.0+	23.5	20.5+	$\checkmark$	
8	Reduce the percentage of adolescents (high school students in grades 9-12) who felt so sad or hopeless every day for 2 weeks or more in a row that they stopped doing some usual activities during the past 12 months	21.8%ª	23%	34.5%	Х	
9	Reduce the mean number of days in the past 30 days adults (age 18 and older) report being mentally unhealthy	3.7	2.9	3.6	Х	
10	Increase the percentage of adolescents (high school students in grades 9-12) with three or more adults (besides their parents) from whom they feel comfortable seeking help	54.2%ª	47%	48.9%	$\checkmark$	
Notes: *Baseline 2010 and current data 2015 unless otherwise noted; DSU = Data statistically unreliable and sup- pressed; +Rate based on fewer than 20 occurrences and statistically unreliable, but not suppressed; ++Modified due to change in data collection methodology; *2009; *2011; *2013; *2006-2010; *2010-2015 For more detailed trend data on any individual indicator, go to http://ibis.dbss.alaska.gov						

**SO** 

For more detailed trend data on any individual indicator, go to http://ibis.dhss.alaska.gov.

Updated 09/0252017




## HEALTHY ALASKANS 2020

A JOINT PROJECT OF THE STATE OF ALASKA DEPARTMENT OF HEALTH AND SOCIAL SERVICES & THE ALASKA NATIVE TRIBAL HEALTH CONSORTIUM

## Healthy Alaskans 2020 Scorecard Regional Edition - Mat-Su

educe the rate of unique substantiated child maltreatment victims per 1,000 children (age 0-17 years) educe the rate of rape per 100,000 population educe the percentage of adolescents (high school students in grades 9-12) who were ever hit, slapped, or physically hurt on purpose of their boyfriend or girlfriend during the past 12 months	Statewide Only Statewide Only	14.4 113.0**	Statewide Only Statewide	
educe the percentage of adolescents (high school students in grades 9-12) who were ever hit, slapped, or physically hurt on purpose	Only	113.0++		
			Only	
	9.0% <sup>c</sup>	8%++	10.4%	Х
educe the alcohol-induced mortality rate per 100,000 population	11.3⁺	15.3	10.5 <sup>+</sup>	$\checkmark$
educe the percentage of adults (age 18 years and older) who report binge drinking in the past 30 days based on the following iteria: 5 or more alcoholic drinks for men; 4 or more alcoholic drinks for women, on one occasion.	24.9%	20%	16.6%	$\checkmark$
educe the percentage of adolescents (high school students in grades 9-12) who report binge drinking in the past 30 days based on e following criteria: 5 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days	24.6%ª	17%	19.4%	Х
educe the unintentional injury mortality rate per 100,000 population	72.7	54.8	64.2	Х
crease the percentage of children age 19-35 months who do receive the ACIP (Advisory Committee on Immunization Practices) commended vaccination series (2013 AICP recommendation: 4 DTaP, 3 Polio, 1MMR, 3 Hib, 3 Hepatitis B, 1 Varicella, 4 PCV)	Statewide Only	75%	Statewide Only	
educe the incidence rate of Chlamydia trachomatis per 100,000 population	486.5	705.2	423.0	$\checkmark$
crease the percentage of rural community housing units with water and sewer services	N/A	87%	N/A	
crease the percentage of the population served by community water systems with optimally fluoridated water	Statewide Only	58%	Statewide Only	
educe the percentage of women delivering live births who have not received prenatal care beginning in the first trimester of pregnancy	22.4%	19%	18.8%	$\checkmark$
educe the rate of preventable hospitalizations per 1,000 adults (hospitalizations that could have been prevented with high quality imary and preventive care) based on the Agency for Healthcare Research and Quality (AHRQ) definition	Statewide Only	6.7	Statewide Only	
educe the rate of adults (age 18 years and older) reporting they could not afford to see a doctor in the past 12 months	18.6%	14%	15.3%	Х
crease the percentage of the population living above the federal poverty level (as defined for AK)	86.9% <sup>d</sup>	90%	86.2% <sup>e</sup>	Х
crease the percentage of 18-24 year olds with a high school diploma or equivalency	75.0% <sup>d</sup>	86%	78.5% <sup>e</sup>	Х
	duce the percentage of adults (age 18 years and older) who report binge drinking in the past 30 days based on the following eria: 5 or more alcoholic drinks for men; 4 or more alcoholic drinks for women, on one occasion. duce the percentage of adolescents (high school students in grades 9-12) who report binge drinking in the past 30 days based on following criteria: 5 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days based on following criteria: 5 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days duce the unintentional injury mortality rate per 100,000 population rease the percentage of children age 19-35 months who do receive the ACIP (Advisory Committee on Immunization Practices) ommended vaccination series (2013 AICP recommendation: 4 DTaP, 3 Polio, 1MMR, 3 Hib, 3 Hepatitis B, 1 Varicella, 4 PCV) duce the incidence rate of Chlamydia trachomatis per 100,000 population rease the percentage of rural community housing units with water and sewer services rease the percentage of the population served by community water systems with optimally fluoridated water duce the percentage of women delivering live births who have not received prenatal care beginning in the first trimester of pregnancy duce the rate of preventable hospitalizations per 1,000 adults (hospitalizations that could have been prevented with high quality nary and preventive care) based on the Agency for Healthcare Research and Quality (AHRQ) definition duce the rate of adults (age 18 years and older) reporting they could not afford to see a doctor in the past 12 months rease the percentage of the population living above the federal poverty level (as defined for AK)	duce the percentage of adults (age 18 years and older) who report binge drinking in the past 30 days based on the following 24.9%   duce the percentage of adolescents (high school students in grades 9-12) who report binge drinking in the past 30 days based on 24.6% <sup>a</sup> duce the percentage of adolescents (high school students in grades 9-12) who report binge drinking in the past 30 days based on 24.6% <sup>a</sup> duce the unintentional injury mortality rate per 100,000 population 72.7   rease the percentage of children age 19-35 months who do receive the ACIP (Advisory Committee on Immunization Practices) Statewide   ommended vaccination series (2013 AICP recommendation: 4 DTaP, 3 Polio, 1MMR, 3 Hib, 3 Hepatitis B, 1 Varicella, 4 PCV) Only   duce the incidence rate of Chlamydia trachomatis per 100,000 population 486.5   rease the percentage of rural community housing units with water and sewer services N/A   rease the percentage of the population served by community water systems with optimally fluoridated water Statewide Only   only duce the rate of preventable hospitalizations per 1,000 adults (hospitalizations that could have been prevented with high quality Statewide Only   duce the rate of adults (age 18 years and older) reporting they could not afford to see a doctor in the past 12 months 18.6%   rease the percentage of the population living above the federal poverty level (as defined for AK) 86.9% <sup>a</sup>	duce the percentage of adults (age 18 years and older) who report binge drinking in the past 30 days based on the following aris: 5 or more alcoholic drinks for men; 4 or more alcoholic drinks for women, on one occasion.24.9%20%duce the percentage of adolescents (high school students in grades 9-12) who report binge drinking in the past 30 days based on following criteria: 5 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days based on following criteria: 5 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days based on the past 30 days based on the past 30 days based on to alcoholic drinks for men; 4 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days based on to alcoholic drinks in a row within a couple of hours, at least once in the past 30 days based on the past 30 days based on the past 30 days based on the past 30 days based on to alcoholic drinks for men; 4 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days based on the following trace the percentage of children age 19-35 months who do receive the ACIP (Advisory Committee on Immunization Practices) ommended vaccination series (2013 AICP recommendation: 4 DTaP, 3 Polio, 1MMR, 3 Hib, 3 Hepatitis B, 1 Varicella, 4 PCV)Statewide Only75%duce the incidence rate of Chlamydia trachomatis per 100,000 population486.5705.2trease the percentage of trual community housing units with water and sewer servicesN/A87%trease the percentage of the population served by community water systems with optimally fluoridated water22.4%19%duce the rate of preventable hospitalizations per 1,000 adults (hospitalizations that could have been prevented with high quality <br< td=""><td>duce the percentage of adults (age 18 years and older) who report binge drinking in the past 30 days based on the following eris. 5 or more alcoholic drinks for wome, on one occasion.24.9%20%16.6%duce the percentage of adolescents (high school students in grades 9-12) who report binge drinking in the past 30 days based on following criteria. 5 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days based on following criteria. 5 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days24.6%*17%19.4%duce the unintentional injury mortality rate per 100,000 population72.754.864.2rease the percentage of children age 19-35 months who do receive the ACIP (Advisory Committee on Immunization Practices) ommended vaccination series (2013 AICP recommendation: 4 DTaP, 3 Polio, 1MMR, 3 Hib, 3 Hepatitis B, 1 Varicella, 4 PCV)Statewide Only75%Statewide Onlyduce the incidence rate of Chamydia trachomatis per 100,000 population486.5705.2423.0423.0rease the percentage of nural community housing units with water and sewer servicesN/A87%N/Aetaes the percentage of fural community housing units with vater and sewer servicesN/A87%Statewide Onlyduce the percentage of preventable hospitalizations per 1,000 adults (hospitalizations that could have been prevented with high quality anary and preventive care) based on the Agency for Healthcare Research and Quality (AHRQ) definition22.4%19%Statewide Onlyduce the rate of preventable hospitalizations per 1,000 adults (hospitalizations that could have been prevented with high quality anary and preventive</td></br<>	duce the percentage of adults (age 18 years and older) who report binge drinking in the past 30 days based on the following eris. 5 or more alcoholic drinks for wome, on one occasion.24.9%20%16.6%duce the percentage of adolescents (high school students in grades 9-12) who report binge drinking in the past 30 days based on following criteria. 5 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days based on following criteria. 5 or more alcoholic drinks in a row within a couple of hours, at least once in the past 30 days24.6%*17%19.4%duce the unintentional injury mortality rate per 100,000 population72.754.864.2rease the percentage of children age 19-35 months who do receive the ACIP (Advisory Committee on Immunization Practices) ommended vaccination series (2013 AICP recommendation: 4 DTaP, 3 Polio, 1MMR, 3 Hib, 3 Hepatitis B, 1 Varicella, 4 PCV)Statewide Only75%Statewide Onlyduce the incidence rate of Chamydia trachomatis per 100,000 population486.5705.2423.0423.0rease the percentage of nural community housing units with water and sewer servicesN/A87%N/Aetaes the percentage of fural community housing units with vater and sewer servicesN/A87%Statewide Onlyduce the percentage of preventable hospitalizations per 1,000 adults (hospitalizations that could have been prevented with high quality anary and preventive care) based on the Agency for Healthcare Research and Quality (AHRQ) definition22.4%19%Statewide Onlyduce the rate of preventable hospitalizations per 1,000 adults (hospitalizations that could have been prevented with high quality anary and preventive

Notes: \*Baseline 2010 and current data 2015 unless otherwise noted; DSU = Data statistically unreliable and suppressed; +Rate based on fewer than 20 occurrences and statistically unreliable, but not suppressed; ++Modified due to change in data collection methodology; <sup>a</sup>2009; <sup>b</sup>2011; <sup>c</sup>2013; <sup>d</sup>2006-2010; <sup>e</sup>2010-2015



No improvement over baseline

Region meets statewide target

 $\checkmark$ 

X Region does not meet statewide target

For more detailed trend data on any individual indicator, go to http://ibis.dhss.alaska.gov.